Astronomy & Astrophysics manuscript no. singles˙arxiv
August 1, 2012

© ESO 2012

arXiv:1207.7229v1 [astro-ph.EP] 31 Jul 2012

Multiple planets or exomoons in *Kepler* hot Jupiter systems with transit timing variations?

R. Szabó[1], Gy. M. Szabó[1,2,3], G. Dálya[4]

[1] Konkoly Observatory, Konkoly-Thege Miklós út 15-17, H-1121 Budapest, Hungary e-mail: `rszabo@konkoly.hu`
[2] Gothard–Lendület Exoplanet Research Team, H-9704 Szombathely, Hungary
[3] Dept. of Exp. Physics & Astronomical Observatory, University of Szeged, H-6720 Szeged, Hungary
[4] Eötvös Loránd University, Pázmány Péter sétány 1/A, H-1117 Budapest, Hungary

Received / Accepted

ABSTRACT

Aims. Hot Jupiters are thought to belong to single-planet systems. Somewhat surprisingly, about one quarter of hot Jupiters were reported to exhibit Transit Timing Variations (TTVs). The aim of this paper is to identify the origin of this observation.
Methods. We present TTV frequencies and amplitudes of hot Jupiters in *Kepler* Q0–6 data with Fourier analysis.
Results. We identified 29 systems with TTV above 4σ confidence, about half of them exhibiting multiple TTV frequencies. Thirteen of these objects (HAT-P-7b, KOI-13, 127, 188, 196, 203, 225, 254, 428, 607, 774, 897, 1176) likely show TTVs due to a systematic observational effect: long cadence data sampling is regularly shifted transit-by-transit, interacting with the transit light curves and introducing a periodic bias. In case of other systems, the activity and rotation of the host star can modulate light curves and explain the observed TTVs. Excluding the systems that were inadequately sampled or showed signs of stellar rotation, we ended up with 16 systems. Nine of them show only one significant period (KOI-131, 186, 256, 882, 897, 1003, 1152, 1448, 1543), others exhibit multiple periods (KOI-823, 977, 1285, 1382, 1452, 1540, 1546). The number of significant periods ranges from 2 (KOI-977, 1382) to 10 (KOI-1546).
Conclusions. The few hot Jupiters with periodic TTVs that we cannot explain with systematics from observation, stellar rotation, activity or inadequate sampling may be good candidates for hot Jupiters in multiple systems or even hosting an exomoon.

Key words. planetary systems – stars: binaries: eclipsing – techniques: photometric

1. Introduction

Transit Timing Variation (TTV) is a major diagnostics of various system parameters of extrasolar planets. In multi-planet systems, planets perturb each other, leading to correlated TTVs of them (Holman et al. 2010, Lissauer et al. 2011). TTVs have also uncovered the presence of further non-transiting planets in solar systems (Ballard et al. 2011, Ford et al. 2012b).

According to our current view, hot Jupiters occur as single planets, since they have been not detected in multi-planet systems. This picture suggests that hot Jupiters occupy unperturbed orbits, hence their orbital motion is Keplerian, and they exhibit strictly periodic transit times. Contrary to this picture, current literature reports a considerable number of hot Jupiters with TTV, which are often periodic (Steffen et al. 2012a, Ford et al. 2012a).

In this work we publish TTV periods, TTV amplitudes and significance levels for hot Jupiter candidates in *Kepler* data. For the analysis, transit times covering Q0–Q6 were taken from Ford et al. 2012a[1] (see Steffen et al. 2012b for more details). We briefly introduce the most exciting systems and discuss the possible source of TTVs for these planets.

2. Data selection and analysis

We have selected all single companions from the second list of *Kepler* planet candidates (Batalha et al. 2012), and filtered the list for planets larger than 6 R_E. Then all systems with incomplete data records, too few observed minima (we required more than 12 transits of a given system for our analysis) or inaccurate orbital periods were neglected. In total, 159 candidates were selected for further analysis. The median planet size and orbital periods were 11.23 R_E and 5.70 days in the sample, the longest orbital period was 54 days. Thus, the sample contains really close-in planets, i.e. they are hot Jupiters. The sample was trimmed further, since some of our targets were reported to be triple systems or blends, thus not hot Jupiters (KOI-190 and KOI-609 Santerne et al. 2012), or the periodic signal was caused by the influence of stellar rotation and/or activity (see in Sec. 4.3). Three systems (KOI-256, KOI-882, KOI-1540) were labelled in Batalha et al. 2012 as showing V-shaped transits. Although planets can cause V-shaped transit light curves, but the probability of being a false positive is much higher for these objects. Despite this, we follow Batalha et al. 2012 in retaining these candidates. Our sample is listed in Table 1. The table contains stellar and planetary parameters taken from the PlanetQuest website[2]. These are compatible with the values from Borucki et al. 2011, Batalha et al. 2012 and the Kepler Input Catalog (Brown et al. 2011). Our table also shows the vetting flag from Borucki et al. 2011, where (2) represents "strong probability candidate, clearly passes tests that were applied", (3) stands for "moderate probability candidate, not all tests cleanly passed, but no definite test failure" and finally (4) means "insufficient follow-up to perform full suit of vetting test."

---

[1] http://www.astro.ufl.edu/~eford/data/kepler/

[2] http://planetquest.jpl.nasa.gov/kepler/



Fig. 1. Illustration of the sub-Nyquist detection of a super-Nyquist modulation in the orbital motion with Fourier analysis. Top panel: one quarter (90 d) long detail from the 600 d long simulations. Dots represent sampling by transits. Bottom panel: Fourier series of the 600 d long data set, extended to beyond the Nyquist frequency. The actual frequency and the sub-Nyquist detections are highlighted. This specific model had an orbital (sampling) and modulation period of 11.21 and 4.76 days (which were taken from the ZIP code and altitude-above-sea-level of the Budapest station of Konkoly Observatory, as uncorrelated random numbers).

Transit times of these systems were taken from the TTV catalog of Ford et al. 2012a and were processed by the standard Fourier analysis with the `MUFRAN` package (Kolláth 1990). Here we wish to illustrate the most serious drawback of the application of the Fourier approach in such an analysis. Once the orbital phase is modulated by a periodic signal, a periodic TTV will be observed. That is, a periodic TTV confidently proves a periodic modulation of the orbital motion. However, the period of the modulation cannot be reconstructed from the TTV if it is shorter than 2 orbital periods. This is because transits undersample such frequent variations, and the observed frequency will be an alias of the super-Nyquist peak. This is illustrated in Fig. 1. In the general case, this can lead to an order-of-magnitude difference between the detected and the actual period. This is unfavourable in the sense that one cannot tell practically anything on the period of the TTV, the detected period will signify only a lower freqeuncy limit for a periodic process. However, the amplitude of the modulation can be properly reconstructed from TTV data (except strictly resonant cases), and the TTV amplitude can be involved into further analysis.

The Fourier-transforms were evaluated individually. The presence of TTVs were characterized by the detected period, the TTV amplitude and the height of the detected peak above the noise level. We considered the planet candidates with at least one peak higher than 4σ above the grass level as systems exhibiting periodic TTVs.



Fig. 2. Calculating the virtual frequency in transit timings, due to the uneven sampling of consecutive transits. P and C are the orbital period and the cadence, respectively, $frac$ means the fractional part.

2.1. Virtual frequencies due to regularly-spaced sampling

We found that in some cases, TTVs exhibited prominent frequencies that can be easily explained by sampling effects. Most of the data are long cadence, and they have a sampling rate of 29.424 minutes. This is in the order of the duration of one whole transit, resulting in 3–6 points belonging to most of the transits. Since the orbital period is not an exact multiple of the cadence rate, the consecutive transits are sampled at different positions, but the lags will evolve very regularly, due to the even sampling rate. The instantly realized sampling can introduce biases in the transit time determination, and because of the regular evolution of the entire process, a virtual variation of transit times may be measured. (A similar feature is seen e.g. in trailed CCD images of asteroids, where the very regular – paced – sampling results in a virtually undulating motion with a sub-pixel amplitude.)

In Fig. 2 we illustrate the derivation of this frequency. Because the cadence of exposures is (obviously) not synchronised to transit mid-times, a systematic lag will develop between the sampling structure of the consecutive transits. The sampling comb will be off one transit later by a time lag of $|P - nC|$, where P and C are the orbital period and the cadence rate, and n is an appropriate integer. We know that $n = [P/C]^{+1}_{-0}$, where [] means the integer part. There are two cases, one when nC is slightly less than P (by less then $P/2$, see this case in Fig. 2), and another one, when nC is slightly larger than P. In the two cases, denoting the fractional part with][brackets, we can write as

$$P - \left[\frac{P}{C}\right]C = C\left(\frac{P}{C} - \left[\frac{P}{C}\right]\right) = C\left]\frac{P}{C}\right[\,, \tag{1}$$



Fig. 3. Sample TTV spectra of systems with monoperiodic modulations. Labeled levels show the confidence limits of peak detections. Spurious peaks are denoted by vertical dashed lines. In case of KOI-1003 after prewhitening with the 277d peak (f=0.003601 d^{-1}), the remaining 108d peak (f=0.00922 d^{-1}) does not reach the 4σ level.

and following the same logic and inverting the sign because the quantity in absolute value brackets is negative

$$\left|P - \left[\frac{P}{C} + 1\right]C\right| = C\left]1 - \frac{P}{C}\right[. \quad (2)$$

Thus, the time lag can be written as $C \cdot s$ if we define $s = \min(]P/C[, 1-]P/C[)$. The $C \cdot s$ time lag is s times one cadence, and after $1/s$ occurrences (i.e transits), the initial cadence-to-transit configuration will be repeated. This means that the virtual peak has P/s period, and hence, s/P frequency. Because s is always less than 0.5 by definition, the virtual peak will always emerge, but will not always be high enough to be detected. In summary, if a transiting system exhibits a TTV period near P/s, this period should be considered to be suspicious, because it may refer to the virtual period of the system.

Among other systems, we found that the most prominent examples of systems with likely virtual TTV periods are KOI-2 (HAT-P-7b), KOI-13, and KOI-127. These systems have a TTV period peak at 21.3, 5.7, and 25.2 days, respectively, while their virtual period is 22.2, 5.7, and 24.3 days. Because we cannot exclude that the detected TTV period is the virtual one, we omitted these objects from our table.

3. Results

Twenty-nine systems out of the 159 were identified showing periodic TTVs, which is 18% of all candidates. Of these targets, 17 candidates show TTV peaks convincingly far from the virtual periods, and also they were not suggested to be blended objects. One more system (KOI-883) was excluded because the TTV period was found to agree with the rotation period of the host star. In Figs. 3. & 4., we plot a sample of Fourier spectra of the found TTV variations.

Surprisingly, 7 of the finally accepted systems exhibited multiple frequency peaks (even 9 and 13 formally above the 4σ level in case of KOI-1452 and KOI-1546, respectively), suggesting that the orbital motion is modulated by several periodic processes. The median period of the detected TTV was 50 days, that means that at least one half of the planet candidates suffer periodic modulations with shorter periods than 50 days. Since the detected TTV period is only an upper limit for the actual period of the modulations, very well may be that the median of the actual periods is much smaller than 50 days, even an order-of-magnitude difference can easily be suspected; but by no ways can the median period exceed 50 days. In some cases (e.g. KOI-1452 and 1540), the major peak is of very low frequency, suggesting a long period process. However, these peaks may also be aliases of rapid oscillations near some resonance with the orbital motion, thus near to the Nyquist frequency multiplied by an even number, and having reflected back near the origin in the sub-Nyquist range. The median amplitude of TTVs is 54 seconds.

We compared system parameters of hot Jupiters with and without TTVs, and found no significant difference in the distributions in the 2-dimensional subspaces of the orbital period, planet size, stellar mass, stellar radius, semi-major axis. One significant difference was found, but very probably due to numerical rather than physical reasons: the orbital period of systems with periodic TTVs is preferentially shorter than that of all hot Jupiters. However, the orbital period is severely biased toward the short period range in the selection process, because a secure detection of any periodic TTV requires many transits, and prefers the short orbital periods.

Since almost all our targets were examined in Steffen et al. 2012b, it is worth comparing our results. We note that our method differs from that of Steffen et al. 2012b: we fit only a single-component sinus at a time, and the definition of the significance level is also different. Instead of taking the uncertainty of the amplitude value, which is a local feature, we compare the amplitude of the signal to the noise computed for the total Fourier-spectrum. We found that the p-values taken from Steffen et al. 2012b are not particularly low for our targets, which means that the TTV values of solitary hot Jupiters are not the highest in the total *Kepler* sample. The SNR values agree for about half of the common objects, while the significance found with our



Fig. 4. Same as Fig. 3., but for systems modulated with multiple periods.

method is usually higher for the rest of our targets. We attribute this difference to the different methods applied.

3.1. Remarks on individual systems

Most of our candidates in Table. 1. are listed in Ford et al. 2011. We only refer to that paper in the table if it provides relevant information in the context of this work.

KOI-131 Santerne et al. 2012 observed this object as part of their large spectroscopic follow-up program of *Kepler* planet candidates. They found a relatively fast rotating exoplanet host star with $v \sin \mathrm{i} = 27 \pm 1\ \mathrm{km\,s^{-1}}$ and place an upper limit of 14.3 M_J for the mass of the companion. The authors note that both the planetary and blend scenarios are compatible with their data. In addition, the object is listed in the *Kepler* eclipsing binary catalog (Prša et al. 2012) with twice the orbital period. This is presumably because of the slightly different depth of the odd and even minima based on the phased light curve on the website[3]. In (Borucki et al. 2011) it got a vetting flag 3. KOI-131 is listed as a planet candidate in Batalha et al. 2012 as well. We conclude that this object is not a strong hot Jupiter candidate.

KOI-186 This Jupiter-sized planet orbits a solar-type star (M=1.06 $M_\odot$). We found two significant periodicities, namely 13.877 and 7.243 days. The later is very close to the presumed stellar rotation period (7.84 d), so we are left with a TTV candidate exhibiting a single periodic variation.

KOI-256 According to Muirhead et al. 2012 the host star is a metal-rich M3 dwarf. Their measurement significantly reduced the size of the planet candidate from 25.34 to 5.60 R_E. The companion shows a single 41.8 d periodicity in its transit timing.

KOI-412 Data from Q5 is missing because the target fell on the failed *Kepler* Module 3. The highest peak in the Fourier spectrum of the transit timing data of this object is located exactly at the Nyquist-frequency, i.e. twice the orbital period of the planet. We can not exclude a signal resulting from a resonant object with 1:2 mean motion resonance in the system. The amplitude of the TTV is uncertain for the same reason.

KOI-822 For similar reasons as in the case of KOI-412, Q5 data are is missing. Strong rotational modulation is seen in the light curve with about half the detected TTV period, as well as its first harmonic, so it may influence the transit timing measurements. Therefore we flagged this star as an uncertain case.

KOI-823 This object fell on the bad module in Q6. It shows complicated, multi-peaked Fourier-spectrum. While some of the peaks are close to peaks seen in the light curve itself (142.9, 59.5, 19.0 d), others are found to be independent of stellar and sampling effects.

KOI-882 Single-periodic TTV candidate. The long period seen in the TTV (41.9 d) is unlikely to be disturbed by other relatively long-period periodicities found in the light curve, the closest one is 37.1 d.

KOI-883 Based on its KIC-parameters, the host star is an early type K dwarf. We found two peaks in the Fourier-spectrum of the transit times of its companion (6.6 d^{-1} and 9.1 d^{-1}). The 6.6 d^{-1} peak is uncomfortably close to the alias caused by the sampling effect, while the latter coincides with the rotation period derived from its light curve presumably caused by the presence of a spot or spots. Thus, this object is no longer a candidate showing periodic TTVs (see Sec. 4.3). We decided the leave it out from Table. 1.

KOI-895 A single-periodic companion around a solar-mass star with 11.4 d TTV period. The stellar rotation, however creates a periodicity of 5.6 d, which is almost exactly half of the TTV period. This renders the main TTV period questionable.

KOI-897 This planetary candidate is slighty larger than Jupiter and orbits a close-to solar mass host star with a period of 2.68 days. There is a single period in its TTV Fourier-spectrum with 82.0 days.

---

[3] http://keplerebs.villanova.edu/

Table 1. Periodic TTV detections above 4-σ level. Periods in italic are ambiguous based on the analysis of the optical light curves of the candidates. The table contains 19 systems, 16 of which shows one or multiple significant TTV periodicities.

KOI	R/R_E	P (d)	M_*	T_{eff}	R_*	TTV per. (d)	Amp. (d)	Sign. lev. (σ)	Vetting flag	Remarks
131.01	9.61	5.0142325	1.13	6244	1.21	114.377216	0.00043	4.2	3	1
186.01	12.35	3.2432603	1.06	5826	0.97	13.877132	0.00027	6.1	2	
256.01	25.34	1.3786789	0.65	3639	0.52	41.755397	0.00031	4.4	3	2
412.01	6.72	4.1470197	1.09	5584	1.17	*8.29*	*>0.00101*	*>4.1*	2	3
822.01	9.79	7.9193704	1	5458	0.74	*24.523629*	*0.00121*	*7.4*	3	4
823.01	7.89	1.028414	1.1	5976	0.96	*142.897970*	*0.00157*	*18.8*	3	
						59.477785	*0.00113*	*13.0*		
						48.562549	0.00090	10.0		
						84.061870	0.00083	9.0		
						343.28870	0.00075	8.0		
						19.033480	0.00061	6.2		
882.01	12.17	1.9568102	0.93	5081	0.55	41.879554	0.00037	6.4	3	2
895.01	10.51	4.4094114	1.04	5436	0.93	*11.436544*	*0.00030*	*5.1*	2	
897.01	12.41	2.0523497	1.07	5734	1.03	81.973932	0.00027	4.7		2
977.01	63.45	1.3537763	0.21	4204	16.48	101.522843	0.00853	5.6	3	5
						20.029644	0.00672	4.0	3	
1003.01	10.84	8.3605703	0.96	5126	0.8	277.700639	0.00207	9.6	3	6
1152.01	15.97	4.7222521	0.58	4069	0.55	*20.868549*	*0.00049*	*9.9*	4	
						11.810977	0.00039	7.5		
1285.01	6.36	0.9374439	0.98	5278	0.83	374.812594	0.00407	65.1	4	7
						106.826193	0.00135	20.3		
						162.786912	0.00104	15.2		
						53.455926	*0.00039*	*4.5*		
						70.229651	*0.00038*	*4.3*		
1382.01	22.64	4.2023359	1.1	5921	1.07	34.449497	0.00102	17.8	4	
						17.397961	0.00068	11.3		
1448.01	24.25	2.4865874	1.08	5658	1.17	11.177805	0.00065	8.4	4	
1452.01	22.39	1.1522169	1.27	6834	1.76	30.290180	0.00148	37.6	4	
						6.308352	0.00126	31.7		
						4.802105	0.00090	22.1		
						74.432453	*0.00088*	*21.6*		
						144.927536	0.00083	20.3		
						44.359668	0.00037	8.1		
						58.309038	*0.00036*	*7.8*		
						192.864031	0.00032	6.7		
						50.584248	0.00030	6.2		
1540.01	31.64	1.2078535	1	5390	0.77	390.015601	0.00043	9.5	4	8
						55.962841	0.00041	8.9		2
						2.426855	0.00041	8.9		
						44.640864	0.00038	8.1		
						2.485046	0.00029	5.8		
						32.99894	*0.00027*	*5.3*		
1543.01	13.69	3.9643337	1.06	5821	0.87	96.99321	0.00031	9.1	4	
1546.01	9.92	0.9175471	0.93	5505	0.86	141.74344	0.00369	77.6	4	
						62.63702	*0.00192*	*39.5*		
						357.90981	*0.00158*	*32.2*		
						74.32181	*0.00104*	*20.6*		
						230.36167	0.00101	20.0		
						107.71219	0.00097	19.1		
						44.161809	0.00064	12.0		
						55.694793	0.00048	8.6		
						33.392326	0.00045	7.9		
						86.625087	0.00038	6.4		
						36.549708	0.00034	5.5		
						10.897400	0.00031	4.9		
						21.113973	0.00029	4.5		

---

[1] Santerne et al. 2012, Prša et al. 2012

[2] V-shaped, Batalha et al. 2012

[3] the peak close to the Nyquist-freq., i.e. 2 P_{orb}

[4] Uncertain: TTV period is close to twice the rotational period. Ford et al. 2011 epoch offset? trend or periodicity

[5] Ford et al. 2011 phase linked variations

[6] Ford et al. 2011 outlier

[7] Ford et al. 2011 spotted star, short duration, epoch offset? trend?

[8] Ford et al. 2011 inaccurate ephemeris in Borucki et al. 2011



Fig. 5. A small part of the *Kepler* light curve of KOI-1152. The transits and eclipses are clearly seen. The eclipses occur at phase 0.34 indicating an eccentric orbit.

KOI-977 The host star of this object has a mass of $0.21M_\odot$ and a radius of $16.48R_\odot$ with an effective temperature of 4204 K according to the Kepler Input Catalog. The nature of this star was corroborated by Muirhead et al. 2012, who clearly identified it as a giant based on near-infrared spectroscopic observations. The radius of the companion is correspondingly large, $63.45R_E$, the largest in our sample. The mass of the host star is obviously too low, most probably a result of the procedure that was optimized to select main-sequence targets for the *Kepler* Mission.

KOI-1003 This is monoperiodic TTV candidate with a long but highly significant period (277 days).

KOI-1152 A larger-than-Jupiter companion orbits an early M1 dwarf (Muirhead et al. 2012). The star shows a 3 day rotation period. Besides the deep transits, the secondary minima (eclipses) are also easily seen. In addition, the orbit of the companion is obviously eccentric, since the eclipses are detected around 0.34 phase instead of 0.50 (see Fig. 5). The lagged secondary eclipse refers to a minimum excentricity of 0.26 (Eq. (2) in Dong et al.2012). The circularization time scale for this system is in the order of $\tau_{circ} \sim$ 120–270 Myr, assuming a planet mass of 1 M_J, a tidal quality factor of 10^6 and other necessary system parameters as derived in KIC and in Batalha et al. 2012. The shorter time scale assumes an excentricity of 0.26, the longer time scale is the $e \sim 0$ limit. Because the circularization time scale is much less than the expected lifetime of an $M_* = 0.58M_\odot$ star, it is reasonable that the system is older than τ_{circ}, and the excentricity is maintained by a dynamical process. Therefore, this system is a good candidate to have an outer companion that perturbs the orbit.

KOI-1285 The optical light curve of this multi-frequency TTV candidate shows peaks close to the 53 and 70 d TTV periodicities, the other 3 long-period variations are not affected.

KOI-1382 This system shows two very clear periodicites (with 34.4 and 17.4 days), close to but not exactly at a 1:2 period ratio. The rotation period of the $1.1M_\odot$ host star is 4.8 days.

KOI-1452 The size of the companion of this star is twice that of the Jupiter. The system exhibits multiple TTV periodicities. The 74.4 and the 58.3 d peaks are close to peaks detected in the Fourier spectrum of the photometric light curve, therefore are ambiguous. The stellar rotation period is estimated to be 1.5 days based on the *Kepler* light curve. The system survived all our criteria, and remains a strong multiperiodic TTV candidate.

KOI-1540 If indeed a planet, this is one of the largest candidate in our vetted sample with almost three Jupiter radius, orbiting a solar-mass star. One of the six significant periods (33 d) is found to be spurious from the light curve analysis.

KOI-1543 KOI-1543 is a monoperiodic TTV candidate with a 3.96 d orbital and a 97.00 d TTV period around a star of $1.06M_\odot$ mass and $0.86R_\odot$ radius.

KOI-1546 This object features the richest spectrum of transit timing variations in our sample. Three frequencies (62.6, 357.9, 74.3 d) are uncertain, because these appear in the light curve as well, another ten peaks reach the 4-σ level.

4. Interpretation

The observation of periodic TTV variations in hot Jupiters has been completely unexpected in the light of the loneliness paradigm, and would imply that one of the following three scenarios acts in systems with hot Jupiters, and affects the transit times:

- systems with transiting hot Jupiters can have non-transiting, massive close-in planets on highly inclined orbits;
- hot Jupiters with significant TTVs host large moons; or
- some other, still unidentified source of periodic TTV is in action.

Here we discuss these possibilities in more details.

4.1. Are there more planets in there?

Hot Jupiters are not necessarily single. If that is the case, periodic TTVs may be tracers of perturbations from additional planets in the system, which are not observed in transits because of the viewing geometry. A known case for a non-single hot Jupiter is in HAT-P-13, hosting a hot Jupiter and a distant massive planet on eccentric orbit, which is not observed in transits (Szabó et al. 2010). In the Batalha et al. 2012 catalog of 2321 planet candidates, there are several examples for hot Jupiters in multiple systems, e.g. KOI-338, a system with two planets, both are hot Jupiters. KOI-94 consists of four transiting candidates, one hot Jupiter, two hot Neptunes and a hot super-Earth, KOI-1241 hosts a hot Jupiter and a hot Neptune; etc.

However, the distribution of hot Jupiter candidates with TTV is very different from the distribution of planets in multi systems. In the left panel of Fig. 6, we plot the apparently single hot Jupiters with crosses and known multiple systems with stars (the left upper part of the region they occupy). We found that very few multiple systems exhibit planets above the line $R/R_E > 5\sqrt[3]{P/day}$. On the other hand, all but one TTV hot Jupiters reside above this line, while they follow the same distribution as non-TTV, (apparently) single hot Jupiters. One could argue that massive, non-transiting companions on an inclined orbit can perturb the transiting planet, but in this case, the planets must be also close-in types to support the median detected TTV period of 50 days. These systems would look very differently from e.g. HAT-P-13.

4.2. Are there exomoons in there?

Another – however, still unconfirmed – explanation of TTVs is the presence of moons, pumping TTVs (Sartoretti and Schenider 1999; Szabó et al. 2006, Simon et al. 2007, Kipping 2009). The expected rate of a TTV can be estimated if we assume that the



Fig. 6. Left panel: test for singleness of TTV hot Jupiters. Stars: multiple systems; crosses: single hot Jupiters; large dots: hot Jupiters with significant TTV. The line shows the upper boundary of the region occupied by known coplanar multi systems. Right: Test for moons. Large dots: hot Jupiters with single TTV periodicity; stars: systems with multiple frequencies taken from Table 1. The line plots the expected TTV amplitudes assuming 1 M_J planet with a 1 M_E moon orbiting at the Hill radius.

satellite has no sign in the light curves, only affects the transit time of the planet. In this case, the TTV full amplitude is $a_s M_s P(\pi a M_p)^{-1}$, where a_s and a is the semi-major axis of the satellite and the planet, M_s and M_p is their mass, respectively, and P is the orbital period (Sartoretti and Schneider, 1999). For an order-of-magnitude estimate of the expected effects, we assigned 1 M_J mass to the planet, 1 M_E mass for the exomoon, and assumed $a_s = a_H$, i.e. the moon orbits at the Hill-radius. With these assumptions, TTV caused by a moon can be calculated as a function of the orbital period of the planet.

In the right panel of Fig. 6, we plot the measured TTV amplitudes for the hot Jupiters. Systems with multiple TTV periods are plotted with different symbols, and they follow the same trend as the monoperiodic systems. The linear model overplots the expected TTV amplitude of a $Jupiter + Earth$ system. We conclude that the distribution of TTV points above an orbital period of 2 days follows a similar trend as the model, but for shorter orbital periods, TTVs significantly exceed this prediction. However, TTV can be underestimated by a factor of 2–5 based on the Sartoretti and Schneider (1999) assumptions especially when the moons are quite large (see Szabó et al. 2006, Simon et al. 2007), and the detected TTVs may be explained with less massive exomoons.

An argument against the exomoon scenario is that a large ($>$ 10%) fraction of hot Jupiters should have distant and massive moons to explain the general appearance of the distribution (right panel of Fig. 6). Besides challenging the origin and prevalence of such systems, the interpretation also faces the question of stability. Hot Jupiters with a moon suffer tidal forces from both the star and from the close-in, massive exomoon. As a consequence, the rotation rate of the planet evolves, and the orbital distance of the exomoon can swing very significantly, either toward spiral-in, or escape, or oscillation, or chaotic behaviour (Barnes & O'Brien 2002). Taken this argument into account, one could raise a reasonable doubt about the assumption of massive exomoons around a significant fraction of hot Jupiters.

4.3. Other scenarios

It is possible that no more objects are necessary to invoke interpreting the observed periodic or quasi-periodic TTVs. Another possible explanation is that the activity of the host stars, i.e. the presence of stellar spots perturb the transit time measurements. An excellent example is Kepler-17b (former KOI-203) which we originally included in our sample. Désert et al. 2011 demonstrated that both the light curve and the O-C of the mid-transit times of this object are modulated by the stellar rotation period and its first harmonic. We also detected the second harmonic in the Fourier spectrum, and therefore discarded this object from further investigations. After scrutinizing the *Kepler* light curve of our target stars, one single periodic target (KOI-883) and about a dozen frequencies in multi-periodic TTV Jupiters were also eliminated because of the suspected indirect effect of the stellar rotation and activity. We note in passing that young, active stars show rotational period less than 10 days, and more aged, quiet stars can exhibit rotation periods up to 30 days. Modulation periods longer than that are unlikely to be caused by activity and rotation.

Conclusions

In this work we selected planetary systems from the *Kepler* sample having only one known hot Jupiter candidates without any other planets. Hot Jupiters are thought to be solitary planets orbiting their host stars. We searched for periodic TTV variations in the sample, a possible sign of exomoons. After eliminating false positives using the results of Santerne et al. 2012 and systems that might show spurious periodic signals due effect of inadequate sampling and contributions from stellar rotation, we ended up with 16 systems. Among these objects 9 show only one significant period (KOI-131, 186, 256, 882, 897, 1003, 1152, 1448, 1543), others show multiple periods (KOI-823, 977, 1285, 1382, 1452, 1540, 1546). The number of significant periods ranges from two (KOI-977, 1382) to ten (KOI-1546). We caution that the detected periods are lower limits, since the sampling is based on the planetary transits in the system, and this can undersample any short-period variations in the Fourier-space. With a median mass of 1.03 $M_{\odot}$ our sample consisting of mainly main-sequence dwarf stars closely follows the distribution of host stellar masses of the Batalha et al. 2012 list (median: 1.05 $M_{\odot}$) with two low-mass stars (KOI-256, 1152) and the evolved KOI-977.

One of our best candidates is KOI-1152, which is an M1 dwarf star hosting a giant planet on a 4.7 d orbit. The eclipses can be detected, and the orbit is obviously eccentric. Since the circularization time is much less than the expected age of the the host star, we hypothesize that the eccentricity and the observed TTV period is maintained by a third body in the system.

Concerning the false positives in our sample, Santerne et al. 2012 recently estimated that the rate of false positives among short-period Jupiters in the *Kepler* sample can be as high as 34.8% ± 6.5%. In the frame of their radial velocity follow-up survey none of our candidates were confirmed as planets, but a few were rejected as we discussed in Sec. 3.1. Previously, Morton & Johnson 2011 estimated a rate which is closer to 5-10 %. Depending on which value is taken, the non-planetary candidates may be between 1 and 6 in our sample, even if brown dwarfs are permitted. More follow-up observations are clearly needed to settle this problem. It is worth noting that 4 of our candidates qualify for a vetting flag of 2, seven of them 3 and the rest (with the highest KOI numbers) have insufficient follow-up observations, therefore have a vetting flag 4 in Borucki et al. 2011.

We discussed three possible scenarios to explain the observed periodic TTVs. If additional planets are present in the systems of these seemingly solitary hot Jupiters, they present a new population of non-transiting, massive, close-in planets, that has not been seen earlier. We emphasize that the TTV hot Jupiters follow the distribution of solitary hot Jupiters on the period-radius diagram.

Although the existence of large (M>1 M_E) exomoons is an attractive possibility, such a scenario needs to address the formation and stability problems of large exomoons around many hot Jupiters. We also eliminated the possibility of the influence of sampling and stellar activity on the observed transit timings.

Longer observations and ground-based follow-ups can help in confirming the nature of the TTV-positive hot Jupiters. With the help of further *Kepler* short cadence observations, or even with accurate transit photometer with large telescopes applying much denser sampling than *Kepler*, sampling effects in long cadence data can be eliminated. With this technique we will be able to confirm that the observed TTVs are not due to effects of regularly spaced sampling with long exposures. Even *Kepler* LC data can help us confirm that the observed TTV is not due to stellar rotation and activity. That is because the appearance of the stellar surface evolves with the activity cycle, thus the effect will be different in different phases of the activity cycle. If the physical nature of the observed hot Jupiter TTVs can be confirmed with appropriate significance, only dynamical explanations survive: additional planets on exotic orbits, or the exomoons.

Acknowledgments

This project has been supported by the Hungarian OTKA Grants K76816, K83790, MB08C 81013 and the “Lendület” Program of the Hungarian Academy of Sciences. RSz and GyMSz were supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences. Eric Ford and Jason Steffen are acknowledged for their help with the electronic table of *Kepler* TTV measurements. We are grateful to the the Kepler Team for making the light curves public and thank all those who worked hard to make this Mission a success.

References

Ballard, S., Fabrycky, D., Fressin, F. et al., 2011, ApJ, 743, 200

Barnes, J. W. & O’Brien, D. P., 2002, ApJ, 575, 1087

Batalha, N. M., Rowe, J. F., Bryson, S. T. et al. 2012, ApJS submitted, arXiv:1202.5852

Bonomo, A. S., Hébrard, G., Santerne, A. et al. 2012, A&A, 538, A96

Borucki, W. J., Koch, D. G., Basri, G. et al. 2011, ApJ, 736, 19

Brown, T. M., Latham, D. W., Everett, M. E., Esquerdo, G. A. 2011, AJ, 142, 112

Désert, J.-M., Charbonneau, D., Demory, B.-O. et al. 2011, ApJS, 197, 14

Dong, S., Katz, B., Socrates, A. 2012, ApJ submitted, arXiv:1201.4399

Ford, E. B., Rowe, J. F., Fabrycky, D. C. et al. 2011, ApJ, 197, 2

Ford, E., Ragozzine, D., Rowe, J. F. et al. 2012, ApJ, accepted, arXiv:1201.1892

Ford, E., Fabrycky, D. C., Steffen, J. H. et al. 2012, ApJ, 750, 113

Holman, M. J., Fabrycky, D. C., Ragozzine, D. et al. 2010, Science, 330, 51

Kipping, D. M., 2009, MNRAS, 396, 1797

Kolláth, Z. 1990, `MUFRAN`, Occ. Tech. Notes of Konkoly Obs., No.1

Lissauer, J. J., Fabrycky, D. C., Ford, E. B. et al. 2011, Nature, 470, 537

Morton, T. D. & Johnson, J. A., 2011, ApJ, 738, 170

Muirhead, P. S., Hamren, K., Schlawin, E. et al. 2012, ApJ, 750, 37

Prša, A., Batalha, N., Slawson, R. W. et al. 2011, AJ, 141, 83

Santerne, A., Díaz, R. F., Moutou, C. et al. 2012, A&A, in press, arXiv:1206.0601

Sartoretti, P. & Schneider, J., 1999, A&AS, 134, 553

Simon, A. E., Szatmáry, K., Szabó, Gy. M., 2007, A&A, 470, 727

Steffen, J. H., Fabrycky, D. C., Ford, E. B. et al. 2012, MNRAS, 421, 2342

Steffen, J. H., Ford, E. B., Rowe, J. F. et al. 2012, ApJ, submitted, arXiv:1201.1873

Szabó, Gy. M., Szatmáry, K., Divéki, Zs., Simon A., 2006, A&A, 450, 395

Szabó, Gy. M., Kiss, L. L., Benkő, J. et al. 2010, A&A, 523, A84